

Forecasting agriculture yield in Nepal using machine learning techniques

Sandeep Dhakal*

Ajaya Dhungana[†]

Abstract

Accurate prediction of agricultural yield is extremely important to ensure food security and cope with the challenges created by climate change and natural disasters. Forecasting agricultural yield is a challenging task due to the complex nature of variables (fertiliser, rainfall, temperature and others) that affect agricultural production. This study employs six supervised machine learning algorithms: Support Vector Machine (SVM), Decision Tree (DT), Random Forest (RF), Multi-Layer Perceptron (MLP), Recurrent Neural Network (RNN), and Convolutional Neural Network (CNN) to build a predictive model using 49 years of historical data (1973-2021) on paddy, wheat, and maize. Model performance was evaluated using Mean Squared Error (MSE), Mean Absolute Error (MAE), Mean Absolute Percentage Error (MAPE), and Root Mean Squared Error (rMSE). Results show that DT and RF models are the most precise with MSE 1% to 5%, MAE 8% to 21%, followed by SVM and CNN. Key predictors of crop yield include area cultivated, capital expenditure, banking expansion, rainfall, temperature, and fertilizers, while irrigation and road network were less significant. The study recommends that farmers prioritize commercial farming, agricultural equipment, and timely availability of fertilizer for application. The Government of Nepal (GoN) should redirect subsidies towards agricultural mechanization, ensure timely supply of fertilizer, and expand banking services in agricultural areas.

Keywords: Agricultural yield, Machine learning, Support vector machine, Random forest, Decision tree, Multilayer perceptron

JEL Classification: C61, C63, Q19

* Corresponding Author. MPhil, Kathmandu University School of Management (KUSOM)

[†] Assistant Director, Securities Board of Nepal

I. INTRODUCTION

Agricultural yield prediction is becoming a more profound issue because of the growing concern of people regarding global food security (OECD, 2013). Even though precise output forecasting is complex and challenging, it is vital for sustainable farming and efficient use of natural resources (Paudel et al., 2021). Accurate estimation of agricultural yield has gained significant importance as it enables policymakers, farmers, agronomists, commodity traders and other stakeholders to make informed decisions and plan strategies to enhance productivity and address potential food shortages. The production of food crops is influenced by a myriad of factors, including crop-specific characteristics, environmental conditions and management decisions (Fischer, 2015). Understanding the interplay between these factors is crucial for developing effective strategies to maximise agricultural yield (Pretty et al., 2006). To address this challenge, researchers have stepped in with advanced technological solutions, particularly machine learning models. Many developed economies have benefited from it. However, countries like Nepal have not been able to take advantage of advanced modelling tools.

The agriculture sector of Nepal, which once dominated the entire economy, is now in decline. The sector contributed 69% to the GDP in 1975, but by 2013/14, this figure dropped to 30.3%, and to 24.1% in 2022/23 (MoF, 2023; World Bank, 2023). The sector grapples with challenges such as inadequate irrigation infrastructure, delayed availability of seeds and fertilizers, subsistence farming, and slow adoption of improved seeds. Moreover, the absence of rural planning exacerbates these issues, leading to unused arable land due to significant youth migration for foreign employment. Over the last decade, the nation observed a modest growth rate of 2.9% in the agriculture sector (MoF, 2023). The labour force dependent on agriculture decreased from 60.4% in 2018 to 50.4% in 2021 (CBS, 2018; MoF, 2023). Nepal has transitioned into a net importer of major crops, with import values of Rs.47.6, Rs.19.6, and Rs.6.32 for paddy, maize, and wheat, respectively, in the fiscal year 2021/22 (MoF, 2023).

Over the past few decades, policymakers introduced various initiatives to support farmers, including the Institutional Development Program, Community Ground Water Irrigation Sector Project, Small Farmer Development Program, and Small Irrigation

Project. Recent efforts involve minimum support prices, 80% subsidy on insurance premiums, digital soil mapping, and an agriculture mechanization project (MoF, 2023). Monetary initiatives, such as priority sector lending, concessional loans, and setting a threshold for agricultural credit as a proportion of total credit, underscore the sector's priority. As of July 2023, total loans disbursed to the agricultural sector amounted to NRS 414.6 billion, constituting 8.5% of the total loan portfolio (NRB, 2023). These interventions underscore the paramount importance of the agricultural sector for policymakers. Given the substantial financial investment by the government, allocating significant resources to bolster this sector, a model for yield prediction becomes not only justifiable but imperative.

Field surveys, crop growth models, remote sensing, statistical models, and various combinations are frequently employed to predict crop yield. With the increasing availability of reliable models and less than adequate precision of existing econometric models, a comprehensive study that objectively assesses the suitability of machine learning models for agricultural yield prediction seems appropriate. The effectiveness of ML in predicting agricultural yield is still unknown for low-income countries whose agricultural ecosystems are not well developed. To the best of the knowledge of the researchers, no published research in Nepal has used ML to predict agricultural yield.

This study is the first of its kind on many counts. First, it is the first study to use machine learning models to predict Nepalese agricultural yield. Nepal is contextually unique in various aspects such as geography, climate, access to technology and scale of farming. Second, this study incorporates several explanatory variables (such as Inflation, Bank Branches and Capital Expenditure) that affect agricultural production but have not been considered in most earlier research. Identification of a suitable forecasting model also benefits all stakeholders. Such studies are valuable to farmers and stakeholders such as traders, researchers, agronomists and policymakers.

The remaining portion of this paper has been presented in the following five sections. Section two presents the review of relevant literature. The research methodology is presented in the third section, the results and discussion of the study are presented in the fourth section, and the study ends with the discussion and conclusion section.

II. LITERATURE REVIEW

A large number of studies have investigated the relationship between agricultural yield and its determinants using different econometric models. Time series and panel data models are the most widely used models (Blanc & Schlenker, 2017; S. R. Singh, 2007). Nguyen et al. (2019) used regression discontinuity to assess the impact of credit policy on rice production in Myanmar. Similarly, Abu and Haruna (2017) and Houensou et al. (2021) used endogenous switching regressions to investigate the impact of access to finance on agricultural commercialization and farm productivity. Weber et al. (2015) used 2SLS for examining crop price impact on agricultural revenue. On the other hand, Headey et al. (2010) and Rada et al. (2011) used data envelopment analysis for assessing agricultural productivity. Studies of D'Agostino and Schlenker (2016), McArthur and McCord (2017), and Schlenker and Roberts (2009) used a fixed effect estimator, and Nguyen Chau and Scrimgeour (2022) and Yitayew et al. (2022) used propensity score matching. Lio and Liu (2006) used Cobb-Douglas production function and Butler and Cornaggia (2011) used DID estimation. Time series and panel data models have a significant contribution to agricultural economic research, but they still possess a limitation, especially in their capacity to predict precisely (Hill et al., 2020; Huang et al., 1998).

With the advancement in statistical computation and estimation methods, several new models based on machine learning and deep learning methods have emerged (Muruganantham et al., 2022). In the last couple of years, machine learning models can be seen used in substantial research in developed economies (Liakos et al., 2018). The concept of machine learning emerged during the 1950s (Samuel, 1959). However, its use as a tool for the estimation of economic variables began quite later. The use of machine learning models for predicting bankruptcy, stock performance, financial distress forecasting and bond rating became quite frequent (Wong & Selvi, 1998) in the 1990s. The utility of the machine learning model in agricultural economics was not common though its use in agriculture can be traced back to Yost et al. (1988). In recent years, research studies have started using machine learning models for agricultural output prediction (Abbaszadeh et al., 2022; Bijanzadeh et al., 2010; Crane-Droesch, 2018; Everingham et al., 2016; M. D. Johnson et al., 2016; Kaul et al., 2005; Koirala et al., 2019; Liu et al., 2001; Pantazi et al., 2016; Wang et al., 2020). More specifically,

Stas et al. (2016) used boosted tree regression and SVM, Liang et al. (2015) used ANN and RF, and Kaul et al. (2005) used ANN and Multiple Linear Regression (MLR). Most of the studies on agricultural yield or yield estimation are based on data from developed economies, especially the US (Ball et al., 2016; Butler & Cornaggia, 2011; Hutchins, 2022; Kukal & Irmak, 2020; Schlenker & Roberts, 2009; Troy et al., 2015; Weber et al., 2015) and other developed economies such as the UK, France, Germany, Italy, Japan and South Korea (Chavas et al., 2019; Corrales et al., 2022; Horie et al., 1992; Landau et al., 1998; Ruß & Brenning, 2010; Yoo et al., 2012).

Existing literature has examined how various factors affect agricultural productivity. However, the application of machine learning models to estimate agricultural yield is still in its early stages. Furthermore, the use of these models in low-income countries like Nepal is uncommon. The capacity of machine learning models to process highly complex data and provide superior fit has made it popular in forecasting agricultural yield. Studies such as Kaul et al. (2005), Koirala et al. (2019), and Pant et al. (2021) noted a mean error of below 10% when predicting yield using machine learning models. Van Klompenburg et al. (2020) noted SVM, DT, RF, and MLP as the most preferred models; therefore, this research has adopted the same to predict the yield of the major crops.

III. RESEARCH METHODOLOGY

Data Collection and Pre-processing

This study tried to incorporate all the possible data; however, data before 1973 was not available for some of the variables. Therefore, the study used data for the period of 1973 to 2021. The study selected only the principal crops (Paddy, Wheat and Maize) for two reasons: first, they occupy a major chunk of the total agricultural production, and second, a complete data set is available for only major crops. The major data sources were Food and Agriculture Organisation (FAO) statistics, Nepal Rastra Bank, the Central Bureau of Statistics of Nepal and the World Bank. The study has altogether used 13 independent variables and three dependent variables. The missing data problem for variables like population, tractors, and road network was encountered. Since the census is conducted once a decade there are only 6 data points from 1971 to 2021. Similarly, road network data collected from economic surveys from 2002 to 2022

was available for only for 14 data points, with a gap of just a year between two data points to a gap of up to 7 years. The variable number of tractors was constructed by taking data from the Department of Transport Management (DOTM) and the FAO. The cumulative number of tractors imported up to the year 2004 was obtained from FAO, and the remaining data was obtained from the DOTM. Linear interpolation and extrapolation methods were used to generate the data for the 49 data points for population, road network and number of tractors. The data was normalised using a standard scalar to scale features to a mean of zero and a standard deviation of one. It ensures that all features contribute equally, speeds up convergence in gradient-based optimization, prevents bias towards features with larger scales, and stabilizes numerical computations. This pre-processing step enhances the accuracy and robustness of the models.

Exploratory Data Analysis (EDA) and Feature Engineering

The study utilized line plots to understand the data patterns. An increasing trend was observed in most of the variables, including crop yield, population, road network, financial deepening, farm size, fertilizer use, irrigation, exchange rate, bank branches, capital expenditure, and per capita income (PCI). Unit root was observed for the majority of the variables, so data was transformed to the first difference. Even after the first difference, per capita income and population were not found stationary, and they were dropped. Similarly, all the variables used for estimation were scaled and normalized to ensure effective learning by the machine. Additionally, during the EDA, the study presented only the mean and standard deviation to describe the data, as the primary focus of this study was on forecasting.

Model Tuning and Validation

The study employed GridSearchCV for tuning SVM, DT, RF, and MLP models, while RandomizedSearchCV was used for RNN and CNN models. Fivefold crossvalidation was conducted for SVM, DT, RF, and MLP, with parameter estimates averaged for accuracy. For CNN and RNN, validation was performed using various train-test splits, ultimately maintaining a 75:25 ratio for final evaluation.

Model Training

The study tried different training test splits and finally used 75:25, realizing its performance as compared to other compositions. This change is now reflected in the methodology section, and the results section discusses its positive impact on model accuracy and reliability. The study utilized machine learning libraries including scikit-learn (version 0.24) and Keras (version 2.4) with TensorFlow backend (version 2.5). The training procedures were executed on a workstation with an Intel i5 processor, 16 GB RAM, and an NVIDIA GeForce 820M GPU to accelerate the training of deep learning models. The training durations varied: traditional machine learning models like SVR, DT, and RF completed training in a few seconds, while more complex models like RNN and CNN required double the time because of their iterative training processes. Standard scaling for features and reshaped input data to fit the requirements of the neural network architectures have been used. These configurations ensured efficient training processes and optimized model performance.

Explanation of variables

The explanatory variables used in this study are farm size, fertilizer, temperature, rainfall, exchange rate, inflation, bank branches, capital expenditure, population and per capita income.

Farm size: Farm size is the size of land in which the crop is cultivated. Farm size is directly proportional to agricultural yield, as large farm sizes benefit from economies of scale, operational efficiency and market power. Studies such as Auffhammer and Carleton (2018) and Cornia (1985) have found positive and significant effects of farm size on agricultural yield, thus motivating us to keep farm size as a predictor variable. In this study, farm size area is in million hectares.

Fertilizer: Fertilizers such as nitrogen, phosphorous, and potassium play a crucial role in growth and development of plants. Fertilizers promote healthy growth by supplying necessary nutrients and by replenishing nutritional deficiencies in the soil. Fertilizers also help to improve pest and disease resistance while also improving soil fertility and sustainability. Studies such as Madzokere et al. (2021), McArthur and McCord (2017), and Wei et al. (2018) have noted positive and significant effects of fertilizer on crop yield. In this study, the aggregate fertilizer data are expressed in thousands of tons.

Temperature and Rainfall: Temperature, precipitation and rainfall have a significant impact on crop productivity. Different crops have specific temperature and rainfall requirements for quality and quantity of yield. Extreme temperatures such as frost and heat waves can damage crops and reduce yield. Similarly, excessive rainfall and soil erosion can impair root health and nutrition availability. Studies such as Auffhammer and Carleton (2018), Chavas et al. (2019), Chen et al. (2022), and D’Agostino and Schlenker (2016) have used temperature and rainfall as predictor variables for agricultural yield.

Irrigation: Efficient irrigation ensures a consistent water supply, promoting optimal crop growth and minimizing water wastage. This contributes to increased agricultural yield and resilience against drought. This study has used land area equipped with irrigation as a proxy for irrigation.

Agricultural Mechanization: Modern machinery streamlines farming tasks, reducing labour and cultivation time. This enhances efficiency, boosts productivity, and improves crop yield and quality. Since data regarding other tools and machinery were unavailable, the number of agricultural tractors has been used as their proxy.

Financial Deepening: Improved financial access enables farmers to invest in advanced technologies and inputs, fostering sustainable practices and resilience against economic uncertainties. Timely credit empowers strategic decision-making for increased agricultural yield. The Private Credit to GDP ratio has been used as a proxy for financial deepening.

Road Network: A well-connected road network facilitates efficient transportation of agricultural produce, reducing post-harvest losses and ensuring timely delivery to markets. This contributes to increased profitability and market integration for farmers. The total road network in kilometres has been used as its proxy.

Inflation: Price level affects input cost, production cost and investment and expansion of the farm. Inflation also affects the purchasing power of consumers, leading to a slash in demand for certain crops and a shift toward lower-cost options. Reduction in demand demotivates producers to produce, thus affecting the overall production

volume. Studies such as Alston et al. (2009), Auffhammer and Carleton (2018), and D. G. Johnson (1980) found a significant impact of price level on agricultural yield.

Population: Population is directly interrelated with food demand, production and agricultural labour supply. A large population creates market opportunities for producers. A large consumer base incentivizes farmers to increase food production to meet the demand. Excessive rise in population may also create challenges for the agriculture sector by increasing pressure on land and by increasing pollution. Studies such as Auffhammer and Carleton (2018), Baffes and Haniotis (2016), and Schneider et al. (2011) have also considered population as a predictor of agricultural yield and price. In this study, population is expressed in millions.

Exchange rate: The exchange rate affects the overall export competitiveness of the country. Low value of domestic currency makes items cheaper for foreign buyers and helps in an increase in demand. Similarly, the exchange rate also influences the cost of imported inputs such as seeds, fertilizer, machinery, and other input items. Obayelu and Salau (2010) and Carter et al. (1990) have found a consequential impact of the exchange rate on agricultural yield. The major trading currency for international trade in Nepal is the USD, ergo, this study has adopted NPR/USD as an explanatory variable.

Bank Branches: A sharp rise in commercial bank and microfinance branches can be seen in Nepal in the last two decades. The number of bank branches has reached 11,569 (NRB, 2023) across the 77 districts of Nepal. Similarly, the loan disbursed by banks to the agricultural sector is Rs.370 billion (NRB, 2023). Bank branches help in agricultural production by providing financial access to the farmers and by supplying them with credit required for necessary investment in irrigation systems and purchased farming equipment, improved seeds and fertilizers, etc. Binswanger et al. (1993) and Khandker and Koolwal (2016) have observed a positive effect of banking expansion on agricultural yield.

Capital expenditure: Government capital expenditure in the form of roads, irrigation, canals, input subsidies, and research and development has a direct effect on the agricultural supply chain and overall productivity. Nepal's government spends around

USD 2 billion per annum for infrastructural development, which is approximately 12 percent of the country's total expenditure. Studies such as Diakosavvas (1990) and Matthew and Mordecai (2016) have noted a sizeable effect of government capital spending on agriculture.

Per Capita Income: Per capita income can have an impact on demand for agricultural products and farmers' access to agricultural inputs. Nepal is a low-income country but its average income per head shows an increasing trend. CBS (2022) reported an increment from USD 260 FY 2000/01 to USD 1246 in FY 2020/21. Lusigi et al. (1998) and Schultz (1956) have also considered per capita income as a probable predictor variable for agricultural yield.

Modelling approach

The general architecture for the prediction of crop yield is presented in Figure 1.




Figure 1: General architecture for crop yield prediction

In this study, four different machine learning models were used in order to predict and back-test the production volume of three major crops. The machine learning data period is segregated into the training and testing periods. A certain portion of historical data is used to train the machine and predict the outcomes in the testing period. But instead of testing for a certain fraction of time, this study has made the prediction for the entire period (both training and testing) so that the accuracy of prediction for both the training and testing periods is visible in graphs. Since there is no fixed rule for splitting the data into training and testing sets, we tried different ratios and found the highest precision with a 75:25 split and reported the same in the data analysis section.

Support Vector Machine (SVM): SVM is a supervised machine learning tool used for the prediction and analysis of complex data using high dimensional space (Vapnik, 1963). Ruß and Kruse (2010) noted SVM as a common machine learning model for predicting crop yield and Pant et al. (2021) have used SVMs for predicting agricultural yield. The goal of SVM is to obtain a function that minimizes the gap between actual and predicted values. The nonlinear model that has been extensively used for modelling non-stationary time series variables and has generated useful results is a support vector. SVR builds linear regression functions as shown below:

$$\hat{y} = w^T x + b \quad (1)$$

Equation (2) shows Vapnik's linear ϵ -Insensitivity loss function.

$$|y - \hat{y}|_{\epsilon} = \begin{cases} 0 & \text{if } |y - \hat{y}| \leq \epsilon \\ |y - f(x_i, w)| - \epsilon, & \text{otherwise} \end{cases} \quad (2)$$

In Equations (1 and 2) \hat{y} represents dependent variables that are paddy, wheat and maize production volume for a given input vector X . The linear regression \hat{y} is estimated by simultaneously minimizing and the sum of the linear ϵ -Insensitivity losses as shown in equation (4). The constant ϵ controls a trade-off between an approximation error and the weight vector.

$$R = \frac{1}{2} \|w\|^2 + c \left(\sum_i^m |y - \hat{y}|_{\epsilon} \right) \quad (3)$$

Minimizing the risk R is equivalent to minimizing the risk shown in Equation 4 under the constraints depicted in Equation 5 – 6. Here, ζ_i and ζ_i^* are surplus variables, one for exceeding the target value by more than ϵ and the other for being more than ϵ below the target.

$$R = \frac{1}{2} \|w\|^2 + c \left(\sum_i^m |y - \zeta_i + \zeta_i^*| \right) \quad (4)$$

$$(w^T X_i + b) - y_i \leq \epsilon + \zeta_i \quad (5)$$

$$y_i - (w^T X_i + b) \leq \epsilon + \zeta_i^* \quad (6)$$

$$\zeta_i \text{ and } \zeta_i^* \geq 0, i = 1, 2, 3, \dots, m \quad (7)$$

Decision Tree (DT): It is a supervised machine learning algorithm used to classify and predict using a non-parametric model. Unlike other estimation tools, a DT does not have a formula. Instead, it builds a tree such as a model with input and its possible consequences. Studies such as Kumar et al. (2020), Pant et al. (2021), and Rajeswari and Suthendran (2019) have used a DT regression model in predicting crop yield.

$$\hat{y} = \sum_i (W_i * f_i(X_i)) \quad (8)$$

In Equation (8), \hat{y} represents the expected value of the dependent variable y for a given input vector X . \sum_i signifies the summation of all the individual nodes. W_i represents the weight assigned to the prediction of the i^{th} node in the tree. Lastly, $f_i(X_i)$ represents the decision rule or function associated with the i^{th} node or leaf in the DT. It takes the input features X_i as input and produces a yield.

Random Forest (RF): RF is an ensemble machine learning tool that uses multiple DTs to predict the targeted variable. Sometimes a single tree is not accurate in predicting the actual value of the dependent variable because of confusion between noise and pattern. Therefore, the creation of n trees increases the chances of accurate prediction where each tree makes an independent prediction, and the final prediction is obtained by assigning weights and averaging them. Studies such as Dang et al. (2021), Everingham et al. (2016), and Jeong et al. (2016) have used a rRF regression model in predicting crop yield.

The formula for the RF regression can be expressed as:

$$\hat{y} = \frac{1}{T} * \sum_i (W_i * f_i(X_i)) \quad (9)$$

Where \hat{y} represents the expected value (or mean) of the dependent variable y for a given input vector X . T denotes the total number of trees in the RF ensemble. \sum_i signifies the summation of all the individual trees in the RF. W_i represents the weight assigned to the prediction of the i^{th} DT in the RF. $f_i(X)$ represents the non-linear prediction function of the i^{th} DT in the RF for the input vector X .

Multilayer perceptron (MLP) regressor: MLP is a type of artificial neural network algorithm that can learn complex relationships between input and yield variables. It

has three layers: input, hidden, and yield. The layers are composed of interconnected neurons flowing from the input to the hidden and the yield layers. Studies such as Gonzalez-Sanchez et al. (2014), Piekutowska et al. (2021), and Ruß and Kruse (2010) have used MLP regressor to predict crop yield. The equation of an MLP regressor can be represented mathematically as follows:

$$\hat{y} = f(W_2 * f(W_1 * X_i + b_1) + b_2) \quad (10)$$

Where, \hat{y} represents dependent variables which are paddy, wheat and maize production volume. X_i represents independent variables which are area cultivated, average temperature, rainfall, fertilizer, BFIs, etc. W_1 and W_2 are the weight matrices that contain the weights for the connections between the input layer and the hidden layer, and between the hidden layer and the yield layer, respectively. b_1 and b_2 are the bias vectors associated with the hidden layer and the yield layer. The function f represents the activation function applied element-wise to the yield of each neuron in the network. During training, the weights (W_1 and W_2) and biases (b_1 and b_2) are adjusted through an optimization algorithm, such as back-propagation, to minimize the difference between the predicted values (\hat{y}) and the actual values of the dependent variable (y). The activation function f introduces non-linearity to the model, allowing it to learn complex relationships between the inputs and the target variable. The architecture for MLP regressor is shown in Figure 2.




Figure 2: MLP architecture for multivariate time series prediction

In the architecture presented in Figure 2, $X_1, X_2, X_3, \dots, X_n$ are input nodes. They represent independent variables which are farm size, fertilizer, rainfall, temperature, bank branches, inflation, exchange rate, inflation, capital expenditure, population and per capita income. \hat{y}_1, \hat{y}_2 , and \hat{y}_3 represent predicted variables which are Paddy, Maize and Wheat. The remaining two are hidden layers.

Recurrent Neural Network (RNN): Let's denote the number of hidden layers as L , where $L = 1, 2, \dots, L$. The equation for input to the first hidden layer is:

$$h_t^l = \text{Activation } W_{ih}^l * x_t + b_{ih}^l + W_{hh}^l * h_{t-1}^l + b_{hh}^l \quad (11)$$

Now, forward pass equation from layer l to hidden Layer $l + 1$ would be;

$$h_t^{l+1} = \text{Activation } W_{ih}^{l+1} * x_t + b_{ih}^{l+1} + W_{hh}^{l+1} * h_{t-1}^{l+1} + b_{hh}^{l+1} \quad (12)$$

And the equation from hidden to yield layer would be;

$$O_t = W_{ho} * h_t^l + b_{ho} \quad (13)$$

Where, h_t^l is the hidden state of hidden layer 1 at time t . W_{ih}^l and W_{hh}^l are the weight matrices for the input-to-hidden and hidden-to-hidden connections for layer l , respectively. b_{ih}^l and b_{hh}^l are the bias vectors for the input-to-hidden and hidden-to-hidden connections for layer 1. W_{ho} is the weight matrix for the hidden-to-yield connections. b_{ho} is the bias vector for the hidden-to-yield connection.

Convolutional Neural Network (CNN): The regression equation for the convolutional neural network is:

$$Y_t = \text{Activation}(W_y * f(h_t) + b_y) \quad (14)$$

Where, Y_t is the predicted yield at time t , h_t is the yield from the convolutional layers, $f(h_t)$ is the operation that flattens or reshapes the yield h_t as needed, W_y is the weight matrix for the regression layer and b_y is the bias vector for the regression layer.

Evaluation Measures

The study conducted by Van Klompenburg et al. (2020) highlighted Mean Absolute Percentage Error (MAPE), Mean Absolute Error (MAE), relative Root Mean Squared Error (rRMSE), Mean Squared Error (MSE), and Coefficient of Determination (R²) as the most preferred performance metrics for the evaluation of machine learning models. Formulas for these evaluation measures are shown in the following equations.

$$MAPE = \frac{1}{n} \sum_{t=1}^n \frac{|y_t - \hat{y}_t|}{|y_t|} * 100 \quad (15)$$

$$MAE = \frac{1}{n} \sum_{t=1}^n \frac{|y_t - \hat{y}_t|}{|y_t|} \quad (16)$$

$$rRMSE = \sqrt{\frac{1}{n} \sum_{t=1}^n \left(\frac{y_t - \hat{y}_t}{y_t} \right)^2} \quad (17)$$

$$MSE = \frac{1}{n} \sum_{t=1}^n (y_t - \hat{y}_t)^2 \quad (18)$$

$$R^2 = 1 - \frac{SSR}{TSS} \quad (19)$$

Where y_t is the actual and \hat{y}_t is the forecasted value.

IV. RESULTS

The results of the study are presented in this section. Table 1 depicts major crop yields, input factors, and other variables. The average paddy yield is 2.54 tons per hectare with a standard deviation of 0.59, indicating a moderate level of variability. Similarly, Maize and Wheat yields have mean values of 1.91 and 1.76 tons per hectare, with standard deviations of 0.44 and 0.57, respectively. The average land area equipped for irrigation is 0.97 million hectares, with a standard deviation of 0.37, highlighting the extent of variability in irrigated land. The average number of tractors is 3.2 thousand units, but the high standard deviation of 4.6 suggests significant variability in tractors across time. The table also presents information on rainfall, temperature, bank branches, exchange rate, inflation, capital expenditure, population, and per capita income.

Table 1: Descriptive Statistics

Variables	Mean	Std. Dev.
Paddy Yield (Tons/Hectare)	2.54	0.59
Maize Yield (Tons/Hectare)	1.91	0.44
Wheat Yield (Tons/Hectare)	1.76	0.57
Paddy (Million Hectare)	1.43	0.11
Maize (Million Hectare)	0.74	0.17
Wheat (Million Hectare)	0.59	0.15
Land area equipped for irrigation (Million Hectare)	0.97	0.37
Tractors (Thousand units)	3.2	4.6
Road network (Km)	18.6	6.9
Fertilizer (1000 tons)	70.14	57.32
Average Temperature (Degree Celsius)	13.99	0.45
Rainfall (mm)	1293.43	147.72
Financial Deepening (Credit/GDP ratio)	30.16	25.05
New Bank Branches	227.08	441.14
Exchange Rate (NRS/USD)	55.03	34.93
Food and Beverage Inflation (%)	8.79	6.13
Capital Expenditure (% of GDP)	7.82	3.35
Population (Million)	21.28	5.22
Per Capita Income (USD)	420.62	367.29

Predictive Performance of Machine Learning Models

Figures 3, 4 and 5 illustrate the predictive performance of the six machine learning models for yield over time. The graphs depict the actual and predicted crop yields across the time-frame of 1973 to 2021. In the figures, Train represents the training period and Test represents the testing period.




Figure 3: Prediction results for paddy yield

Figure 3 shows predictions of the first difference in paddy yield. During the training period, all models (SVM, DT, RF, MLP, RNN, CNN) performed well, closely following the actual values. However, in the testing period, the models' performance varies. SVM shows reasonable predictions with some deviations. DT struggles significantly, while RF and MLP exhibit noticeable divergences. RNN also shows reduced accuracy, and CNN, at though better than the others, still has periods of deviation. Overall, these models face challenges in maintaining accuracy during the testing period, highlighting potential overfitting or limitations in capturing yield patterns.




Figure 4: Prediction results for wheat yield

In Figure 4, the DT and the RF models show good performance during training but perform poorly during testing. Similarly, the SVM and the MLP models exhibit accurate predictions in the training phase but lack predictive power in the testing phase. The RNN model displays significant fluctuations and inaccuracies, particularly in the testing period. Conversely, the CNN model consistently delivers satisfactory performance across both the training and testing periods, despite some errors in the Test phase.




Figure 5: Prediction results for maize yield

In Figure 5, the SVM model provides decent predictions in the training phase but loses accuracy in the testing phase. The MLP model displays significant errors during the testing period, indicating poor generalization. The RNN model exhibits considerable fluctuations and inaccuracies, especially in the testing period. The CNN model, despite some errors in the Test phase, demonstrates relatively consistent performance across both training and the testing periods. DT and RF perform relatively better in the testing period as compared to other models.

Table 2: Performance Metrics

Product	Model	MAE		MSE		RMSE		MAPE	
		Train	Test	Train	Test	Train	Test	Train	Test
Paddy	SVM	0.10	0.19	0.02	0.04	0.13	0.21	3243.6	144.5
	DT	0.00	0.21	0.00	0.05	0.00	0.23	0.0	193.4
	RF	0.07	0.17	0.01	0.03	0.09	0.19	414.1	103.1
	MLP	0.08	0.85	0.01	0.91	0.11	0.95	2428.2	1518.2
	RNN	0.17	0.23	0.05	0.08	0.23	0.28	6182.0	346.1
	CNN	0.10	0.43	0.02	0.25	0.15	0.50	827.2	731.3
Wheat	SVM	0.06	0.16	0.00	0.03	0.07	0.17	134.4	94.0
	DT	0.00	0.21	0.00	0.05	0.00	0.23	0.0	137.5
	RF	0.03	0.15	0.00	0.03	0.04	0.17	60.2	96.4
	MLP	0.05	0.65	0.00	0.58	0.06	0.76	134.1	411.6
	RNN	0.07	1.35	0.01	2.63	0.09	1.62	136.4	915.9
	CNN	0.05	0.18	0.00	0.05	0.06	0.22	156.2	114.4
Maize	SVM	0.07	0.11	0.01	0.01	0.09	0.12	154.2	101.9
	DT	0.00	0.10	0.00	0.01	0.00	0.11	0.0	94.1
	RF	0.04	0.08	0.00	0.01	0.05	0.10	54.4	78.1
	MLP	0.05	0.17	0.01	0.05	0.08	0.23	103.4	199.4
	RNN	0.15	0.47	0.04	0.32	0.19	0.56	474.8	575.7
	CNN	0.06	0.16	0.01	0.04	0.09	0.19	130.0	159.0

Table 2 shows performance metrics of different machine learning models (SVM, DT, RF, MLP, RNN, CNN) for predicting Paddy, Wheat, and Maize yields. For Paddy, RF performs best with a Test MAE of 0.17 and Test MAPE of 103.1, while SVM and CNN show moderate errors and MLP and RNN struggle with higher errors. For Wheat, RF again excels with a Test MAE of 0.15 and Test MAPE of 96.4, whereas SVM and CNN perform moderately, and MLP and RNN have significant challenges. For Maize, RF and DT are the most effective, with RF showing the lowest Test MAE of 0.08 and Test MAPE of 78.1. SVM also performs well, while MLP and RNN exhibit higher errors. Tree-based models likely perform well due to their ability to handle non-linear relationships and interactions within the data, making them well-suited for agricultural yield prediction. In contrast, the RNN and MLP models may struggle with capturing the complex temporal and spatial dependencies inherent in crop yield data. Overall,

RF is the most promising model across all crops, while RNN and MLP struggle and CNN shows variable performance depending on the crop, suggesting it may be better suited for specific patterns or dependencies.

Feature Importance

Since a superior fit was found only for RF and DT during the testing period, feature importance was calculated using these two models. For each dependent variable, the data was split into training and testing sets after scaling, and feature importances were extracted.

Table 3: Feature Importance

Features	Paddy		Maize		Wheat	
	RF	DT	RF	DT	RF	DT
Capital Expenditure	0.381	0.182	0.156	0.120	0.156	0.080
Banking Expansion	0.183	0.077	0.159	0.077	0.09	0.09
Area Cultivated	0.111	0.148	0.21	0.256	0.13	0.14
Food and Beverage Inflation	0.04	0.11	0.19	0.0665	0.174	0.189
Agricultural Mechanization	0.07	0.09	0.068	0.113	0.14	0.130
Temperature	0.04	0.075	0.06	0.054	0.064	0.12
Rainfall	0.11	0.185	0.054	0.183	0.140	0.098
Fertilizers	0.05	0.087	0.04	0.085	0.044	0.13

Table 3 indicates the significance of each variable in predicting agricultural productivity using both RF and DT models. For Paddy, Capital Expenditure emerges as the most crucial factor in the RF model with a high importance value of 0.381, followed by Banking Expansion and Area Cultivated. The DT model also places importance on Capital Expenditure (0.182) but highlights Area Cultivated (0.148) as more significant. For Maize, the Area Cultivated is the most influential variable in both RF (0.21) and DT (0.256) models, with Capital Expenditure and Banking Expansion also showing significant importance. In the case of Wheat, Food and Beverage Inflation stands out as the most important feature in both RF (0.174) and DT (0.189) models. Capital Expenditure and Agricultural Mechanization are also critical factors in the RF model, while the DT model assigns considerable importance to Area Cultivated and

Agricultural Mechanization. Interestingly, variables like Temperature and Fertilizers have lower importance across both models for all crops, indicating a lesser impact on productivity predictions.

V. DISCUSSION AND CONCLUSION

Predicting crop yield is challenging because of several factors affecting their productivity both directly and indirectly. This study builds a predictive model using three major crops and fourteen key indicators driving their productivity. Six supervised machine learning algorithms, SVM, DT, RF MLP, RNN and CNN, were used. Experiments were conducted using 49 years of historical data from 1973 to 2021 on three principal crops — paddy, wheat, and maize; and the model performance was measured using MSE, MAE, MAPE, and rMSE. DT and RF have been found to be the most precise, followed by SVM and CNN. The outcomes are robust across crop types. Additionally, variables such as area cultivated, capital expenditure, banking expansion, rainfall, temperature and fertilizers have been found to be the most important features for predicting crop yield. Surprisingly, feature importance did not show irrigation, road network, and financial deepening as key drivers of yield. DT observed a MAE of 21%, for paddy and wheat and 10% for maize. Similarly, RF found an MAE of 17% for paddy, 15% for wheat, and 8% for maize. Similarly, MSE is lowest for DT and RF for all crop types. Different from our findings, Kung et al. (2016) observed an error of only 1.3% in general and identified agricultural, meteorological and harvest data as key features. Similarly, Pantazi et al. (2016) obtained an accuracy of 81.6% while estimating wheat yield within field variation. Su et al. (2017) modelled to predict rice yield with significantly low rMSE and found soil quality and surface weather as major features. Amaratunga et al. (2020) observed the mean square error from 2% to as high as 38% in ANN. P. K. Singh et al. (2022) observed relatively low MAPE and MSE when applied to SVM. According to Kang et al. (2020) XGBoost algorithm outperforms other algorithms, both in accuracy and stability, while deep neural networks such as LSTM and CNN are not advantageous. As per Grinsztajn et al. (2022) tree based models like RF and DT perform better due to their ability to handle non-linear relationships and their robustness to outliers.

Research indicates that area cultivated, use of fertilizers, and agricultural

mechanization are key drivers of crop yield. To improve yields, farmers should prioritize allocating more resources towards commercial farming practices, investing in essential agricultural equipment such as tractors, and ensuring the timely and adequate application of fertilizers. Additionally, the GoN has been investing billions of rupees in agricultural subsidies. To further boost crop yields, it is recommended that GoN redirect subsidies towards the acquisition of agricultural machinery, guarantee timely and sufficient supply of fertilizers to farmers, and establish more bank branches in agricultural areas to facilitate easier access to financial services and support. By focusing on these areas, both farmers and the government can work together to directly and indirectly enhance crop yields, leading to increased agricultural productivity and food security. Based on the findings of this study, diverting resources to increasing cultivation area, capital expenditure, banking expansion, climatic conditions, and fertilizers is likely to help increase yield. Similarly, instead of applying only a linear prediction model, it is recommended to use a tree-based model like DT or RF for a superior fit.

Regarding the limitations, this study could not incorporate key indicators like agricultural tools, fertilizer type, and policy variables, and the study used yearly data of only 49 years due to the unavailability of data. Future research in this field could incorporate variables like irrigation, agricultural tools, fertilizer type, and other socio-economic and policy variables for improved prediction capacity. Future research can also utilize district-wise panel data as both temporal and spatial data would provide sizeable data for the model to train itself and provide a superior fit.

For future research, one of the most challenging aspects will be obtaining crop-specific data. The reliability of predictions will significantly improve with detailed crop-wise input data such as irrigation availability, fertilizer use, seed quality, labour, and average farm size. Variables like rainfall, temperature, and prices, which affect yield, are inherently random and exhibit stochastic properties. Integrating stochastic models like Vasicek, Heston, or Stochastic Block Allocation Regression (SBAR) alongside crop-specific variables presents both opportunities and challenges for future research.

REFERENCES

- Abbaszadeh, P., Gavahi, K., Alipour, A., Deb, P., & Moradkhani, H. (2022). Bayesian multi-modeling of deep neural nets for probabilistic crop yield prediction. *Agricultural and Forest Meteorology*, 314, 108773. <https://doi.org/10.1016/j.agrformet.2021.108773>
- Abu, B. M., & Haruna, I. (2017). Financial inclusion and agricultural commercialization in Ghana: An empirical investigation. *Agricultural Finance Review*, 77 (4), 524–544. <https://doi.org/10.1108/AFR-02-2017-0007>
- Alston, J. M., Beddow, J. M., & Pardey, P. G. (2009). Agricultural research, productivity, and food prices in the long run. *Science*, 325 (5945), 1209–1210. <https://doi.org/10.1126/science.1170451>
- Amaratunga, V., Wickramasinghe, L., Perera, A., Jayasinghe, J., & Rathnayake, U. (2020). Artificial neural network to estimate the paddy yield prediction using climatic data. *Mathematical Problems in Engineering*, 2020 (1), 8627824.
- Auffhammer, M., & Carleton, T. A. (2018). Regional crop diversity and weather shocks in India. *Asian Development Review*, 35 (2), 113–130. <https://doi.org/10.1162/adev.a.00116>
- Baffes, J., & Hanjotis, T. (2016). What explains agricultural price movements? *Journal of Agricultural Economics*, 67 (3), 706–721. <https://doi.org/10.1111/1477-9552.12172>
- Ball, E. V., Wang, S. L., Nehring, R., & Mosheim, R. (2016). Productivity and economic growth in US agriculture: A new look. *Applied Economic Perspectives and Policy*, 38 (1), 30–49. <https://doi.org/10.1093/aepp/ppv031>
- Bijanzadeh, E., Emam, Y., & Ebrahimie, E. (2010). Determining the most important features contributing to wheat grain yield using supervised feature selection model. *Australian Journal of crop science*, 4 (6), 402–407.
- Binswanger, H. P., Khandker, S. R., & Rosenzweig, M. R. (1993). How infrastructure and financial institutions affect agricultural output and investment in India. *Journal of development Economics*, 41 (2), 337–366. [https://doi.org/10.1016/0304-3878\(93\)90062-R](https://doi.org/10.1016/0304-3878(93)90062-R)

- Blanc, E., & Schlenker, W. (2017). The use of panel models in assessments of climate impacts on agriculture. *Review of Environmental Economics and Policy*. <https://doi.org/10.1093/reep/rex016>
- Butler, A. W., & Cornaggia, J. (2011). Does access to external finance improve productivity? Evidence from a natural experiment. *Journal of Financial Economics*, 99 (1), 184–203. <https://doi.org/10.1016/j.jfineco.2010.08.009>
- Carter, C. A., Gray, R. S., & Furtan, W. (1990). Exchange rate effects on inputs and outputs in Canadian agriculture. *American Journal of Agricultural Economics*, 72 (3), 738–743. <https://doi.org/10.2307/1243047>
- CBS. (2018). *Report on the Nepal Labour Force Survey, 2017/18* (tech. rep.) (Accessed on 13th December 2023). National Planning Commission, National Bureau of Statistics. <https://cbs.gov.np/wpcontent/uploads/2019/05/Nepal-Labour-Force-Survey-2017-18-Report.pdf>
- CBS. (2022). *National Accounts FY 2021/22* (National Report). Central Bureau Statics, Nepal. Kathmandu.
- Chavas, J.-P., Di Falco, S., Adinolfi, F., & Capitanio, F. (2019). Weather effects and their long-term impact on the distribution of agricultural yields: Evidence from Italy. *European Review of Agricultural Economics*, 46 (1), 29–51. <https://doi.org/10.1093/erae/jby019>
- Chen, B., Dennis, E. J., & Featherstone, A. (2022). Weather impacts the agricultural production efficiency of wheat: The emerging role of precipitation shocks. *Journal of Agricultural and Resource Economics*, 47 (3), 544–562.
- Cornia, G. A. (1985). Farm size, land yields and the agricultural production function: An analysis for fifteen developing countries. *World development*, 13 (4), 513–534. [https://doi.org/10.1016/0305-750X\(85\)90054-3](https://doi.org/10.1016/0305-750X(85)90054-3)
- Corrales, D. C., Schoving, C., Raynal, H., Debaeke, P., Journet, E.-P., & Constantin, J. (2022). A surrogate model based on feature selection techniques and regression learners to improve soybean yield prediction in southern France. *Computers and Electronics in Agriculture*, 192, 106578. <https://doi.org/10.1016/j.compag.2021.106578>

- Crane-Droesch, A. (2018). Machine learning methods for crop yield prediction and climate change impact assessment in agriculture. *Environmental Research Letters*, 13 (11), 114003. <https://doi.org/10.1088/1748-9326/aae159>
- D’Agostino, A. L., & Schlenker, W. (2016). Recent weather fluctuations and agricultural yields: Implications for climate change. *Agricultural economics*, 47 (S1), 159–171. <https://doi.org/10.1111/agec.12315>
- Dang, C., Liu, Y., Yue, H., Qian, J., & Zhu, R. (2021). Autumn crop yield prediction using data-driven approaches:-support vector machines, random forest, and deep neural network methods. *Canadian journal of remote sensing*, 47 (2), 162–181. <https://doi.org/10.1080/07038992.2020.1833186>
- Diakosavvas, D. (1990). Government expenditure on agriculture and agricultural performance in developing countries: An empirical evaluation. *Journal of Agricultural Economics*, 41 (3), 381–389. <https://doi.org/10.1111/j.14779552.1990.tb00654.x>
- Everingham, Y., Sexton, J., Skocaj, D., & Inman-Bamber, G. (2016). Accurate prediction of sugarcane yield using a random forest algorithm. *Agronomy for sustainable development*, 36, 1–9. <https://doi.org/10.1007/s13593-016-0364-z>
- Fischer, R. (2015). Definitions and determination of crop yield, yield gaps, and of rates of change. *Field Crops Research*, 182, 9–18. <https://doi.org/10.1016/j.fcr.2014.12.006>
- Gonzalez-Sanchez, A., Frausto-Solis, J., & Ojeda-Bustamante, W. (2014). Predictive ability of machine learning methods for massive crop yield prediction. *Spanish Journal of Agricultural Research*, 12 (2), 313–328. <https://doi.org/10.5424/sjar/2014122-4439>
- Grinsztajn, L., Oyallon, E., & Varoquaux, G. (2022). Why do tree-based models still outperform deep learning on typical tabular data? *Advances in neural information processing systems*, 35, 507–520.
- Headey, D., Alauddin, M., & Rao, D. P. (2010). Explaining agricultural productivity growth: An international perspective. *Agricultural Economics*, 41 (1), 1–14. <https://doi.org/10.1111/j.1574-0862.2009.00420.x>

- Hill, T. D., Davis, A. P., Roos, J. M., & French, M. T. (2020). Limitations of fixed-effects models for panel data. *Sociological Perspectives*, 63 (3), 357–369. <https://doi.org/10.1177/0731121419863785>
- Horie, T., Yajima, M., & Nakagawa, H. (1992). Yield forecasting. *Agricultural systems*, 40 (1-3), 211–236. [https://doi.org/10.1016/0308-521X\(92\)90022-G](https://doi.org/10.1016/0308-521X(92)90022-G)
- Houensou, D. A., Goudjo, G. G., & Senou, M. M. (2021). Access to finance and difference in family farm productivity in Benin: Evidence from small farms. *Scientific African*, 13, 1–12. <https://doi.org/10.1016/j.sciaf.2021.e00940>
- Huang, N. E., Shen, Z., Long, S. R., Wu, M. C., Shih, H. H., Zheng, Q., Yen, N.-C., Tung, C. C., & Liu, H. H. (1998). The empirical mode decomposition and the hilbert spectrum for nonlinear and non-stationary time series analysis. *Proceedings of the Royal Society of London. Series A: mathematical, physical and engineering sciences*, 454 (1971), 903–995. <https://doi.org/10.1098/rspa.1998.0193>
- Hutchins, J. (2022). The US farm credit system and agricultural development: Evidence from an early expansion, 1920–1940. *American Journal of Agricultural Economics*, 105 (1), 3–26. <https://doi.org/10.1111/ajae.12290>
- Jeong, J. H., Resop, J. P., Mueller, N. D., Fleisher, D. H., Yun, K., Butler, E. E., Timlin, D. J., Shim, K.-M., Gerber, J. S., & Reddy, V. R. (2016). Random forests for global and regional crop yield predictions. *PloS one*, 11 (6), e0156571. <https://doi.org/10.1371/journal.pone.0156571>
- Johnson, D. G. (1980). Inflation, agricultural output, and productivity. *American Journal of Agricultural Economics*, 62 (5), 917–923. <https://doi.org/10.2307/1240284>
- Johnson, M. D., Hsieh, W. W., Cannon, A. J., Davidson, A., & Bédard, F. (2016). Crop yield forecasting on the Canadian Prairies by remotely sensed vegetation indices and machine learning methods. *Agricultural and forest meteorology*, 218, 74–84. <https://doi.org/10.1016/j.agrformet.2015.11.003>
- Kang, Y., Ozdogan, M., Zhu, X., Ye, Z., Hain, C., & Anderson, M. (2020). Comparative assessment of environmental variables and machine learning algorithms for maize yield prediction in the us midwest. *Environmental Research Letters*, 15 (6), 064005.

- Kaul, M., Hill, R. L., & Walthall, C. (2005). Artificial neural networks for corn and soybean yield prediction. *Agricultural Systems*, 85 (1), 1–18. <https://doi.org/10.1016/j.agsy.2004.07.009>
- Khandker, S. R., & Koolwal, G. B. (2016). How has microcredit supported agriculture? Evidence using panel data from Bangladesh. *Agricultural Economics*, 47 (2), 157–168. <https://doi.org/10.1111/agec.12185>
- Koirala, A., Walsh, K. B., Wang, Z., & McCarthy, C. (2019). Deep learning—method overview and review of use for fruit detection and yield estimation. *Computers and electronics in agriculture*, 162, 219–234. <https://doi.org/10.1016/j.compag.2019.04.017>
- Kukul, M., & Irmak, S. (2020). Impact of irrigation on interannual variability in united states agricultural productivity. *Agricultural Water Management*, 234, 106141. <https://doi.org/10.1016/j.agwat.2020.106141>
- Kumar, Y. J. N., Spandana, V., Vaishnavi, V., Neha, K., & Devi, V. (2020). Supervised machine learning approach for crop yield prediction in agriculture sector. *2020 5th International Conference on Communication and Electronics Systems (ICCES)*, 736–741. <https://doi.org/10.1109/ICCES48766.2020.9137868>
- Kung, H.-Y., Kuo, T.-H., Chen, C.-H., & Tsai, P.-Y. (2016). Accuracy analysis mechanism for agriculture data using the ensemble neural network method. *Sustainability*, 8 (8), 735.
- Landau, S., Mitchell, R., Barnett, V., Colls, J., Craigon, J., Moore, K., & Payne, R. (1998). Testing winter wheat simulation models’ predictions against observed UK grain yields. *Agricultural and Forest Meteorology*, 89 (2), 85–99. [https://doi.org/10.1016/S0168-1923\(97\)00069-5](https://doi.org/10.1016/S0168-1923(97)00069-5)
- Liakos, K. G., Busato, P., Moshou, D., Pearson, S., & Bochtis, D. (2018). Machine learning in agriculture: A review. *Sensors*, 18 (8), 2674. <https://doi.org/10.3390/s18082674>
- Liang, L., Di, L., Zhang, L., Deng, M., Qin, Z., Zhao, S., & Lin, H. (2015). Estimation of crop lai using hyperspectral vegetation indices and a hybrid inversion method. *Remote Sensing of Environment*, 165, 123–134. <https://doi.org/10.1016/j.rse.2015.04.032>

- Lio, M., & Liu, M.-C. (2006). ICT and agricultural productivity: evidence from crosscountry data. *Agricultural Economics*, 34 (3), 221–228. <https://doi.org/10.1111/j.1574-0864.2006.00120.x>
- Liu, J., Goering, C., & Tian, L. (2001). A neural network for setting target corn yields. *Transactions of the ASAE*, 44 (3), 705. <https://doi.org/10.13031/2013.6097>
- Lusigi, A., Piesse, J., & Thirtle, C. (1998). Convergence of per capita incomes and agricultural productivity in Africa. *Journal of International Development: The Journal of the Development Studies Association*, 10 (1), 105–115.
- Madzokere, T. C., Murombo, L., & Chiririwa, H. (2021). Nano-based slow releasing fertilizers for enhanced agricultural productivity. *Materials Today: Proceedings*, 45, 3709–3715. <https://doi.org/10.1016/j.matpr.2020.12.674>
- Matthew, A., & Mordecai, B. D. (2016). The impact of public agricultural expenditure on agricultural output in Nigeria (1981-2014). *Asian Journal of Agricultural Extension, Economics & Sociology*, 11 (2), 1–10. <https://doi.org/10.9734/AJAEES/2016/25491>
- McArthur, J. W., & McCord, G. C. (2017). Fertilizing growth: Agricultural inputs and their effects in economic development. *Journal of development economics*, 127, 133–152. <https://doi.org/10.1016/j.jdeveco.2017.02.007>
- MoF. (2023). *Economic Survey* (Annual Report). Government of Nepal, Ministry of Finance. <https://www.mof.gov.np/uploads/document/file/1687415151Economic%20Survey%207980.pdf>
- Muruganantham, P., Wibowo, S., Grandhi, S., Samrat, N. H., & Islam, N. (2022). A systematic literature review on crop yield prediction with deep learning and remote sensing. *Remote Sensing*, 14 (9), 1990.
- Nguyen, H.-T.-M., Sparrow, R., & Aung, N. (2019). The Impact of Credit Policy on Rice Production in Myanmar. *Journal of Agricultural Economics*. <https://doi.org/10.1111/1477-9552.12299>
- Nguyen Chau, T., & Scrimgeour, F. (2022). Productivity impacts of hybrid rice seeds in Vietnam. *Journal of Agricultural Economics*, 73 (2), 414–429. <https://doi.org/10.1111/1477-9552.12458>

- NRB. (2023). *Banking and financial statistics* (Monthly Report). Nepal Rashtra Bank. [https://www.nrb.org.np/contents/uploads/2023/09/Saun 2080 Publish-3.pdf](https://www.nrb.org.np/contents/uploads/2023/09/Saun%202080Publish-3.pdf)
- Obayelu, A. E., & Salau, A. S. (2010). Agricultural response to prices and exchange rate in nigeria: Application of co-integration and vector error correction model (vecm). *Journal of agricultural sciences*, 1 (2), 73–81. <https://doi.org/10.1080/09766898.2010.11884656>
- OECD. (2013). *Global food security: Challenges for the food and agricultural system*.
- Pant, J., Pant, R., Singh, M. K., Singh, D. P., & Pant, H. (2021). Analysis of agricultural crop yield prediction using statistical techniques of machine learning. *Materials Today: Proceedings*, 46, 10922–10926. <https://doi.org/10.1016/j.matpr.2021.01.948>
- Pantazi, X. E., Moshou, D., Alexandridis, T., Whetton, R. L., & Mouazen, A. M. (2016). Wheat yield prediction using machine learning and advanced sensing techniques. *Computers and electronics in agriculture*, 121, 57–65. <https://doi.org/10.1016/j.compag.2015.11.018>
- Paudel, D., Boogaard, H., de Wit, A., Janssen, S., Osinga, S., Pylaniadis, C., & Athanasiadis, I. N. (2021). Machine learning for large-scale crop yield forecasting. *Agricultural Systems*, 187, 1–13. <https://doi.org/10.1016/j.agsy.2020.103016>
- Piekutowska, M., Niedba-la, G., Piskier, T., Lenartowicz, T., Pilarski, K., Wojciechowski, T., Pilarska, A., & Czechowska-Kosacka, A. (2021). The application of multiple linear regression and artificial neural network models for yield prediction of very early potato cultivars before harvest. *Agronomy*, 11 (5), 885. <https://doi.org/10.3390/agronomy11050885>
- Pretty, J. N., Noble, A. D., Bossio, D., Dixon, J., Hine, R. E., Penning de Vries, F. W., & Morison, J. I. (2006). Resource-conserving agriculture increases yields in developing countries. *Environmental Science and Technology*, 40 (4), 1114–1119. <https://doi.org/10.1021/es051670d>
- Rada, N. E., Buccola, S. T., & Fuglie, K. O. (2011). Government policy and agricultural productivity in Indonesia. *American journal of agricultural economics*, 93 (3), 867–884. <https://doi.org/10.1093/ajae/aar004>

- Rajeswari, S., & Suthendran, K. (2019). Advanced Decision Tree (ADT) classification model for agricultural data analysis on cloud. *Computers and Electronics in Agriculture*, 156, 530–539. <https://doi.org/10.1016/j.compag.2018.12.013>
- Ruß, G., & Brenning, A. (2010). Spatial variable importance assessment for yield prediction in precision agriculture. *Advances in Intelligent Data Analysis IX: 9th International Symposium, IDA 2010, Tucson, AZ, USA, May 19–21, 2010. Proceedings 9*, 184–195.
- Ruß, G., & Kruse, R. (2010). Regression models for spatial data: An example from precision agriculture. *Advances in Data Mining. Applications and Theoretical Aspects: 10th Industrial Conference, ICDM 2010, Berlin, Germany, July 12–14, 2010. Proceedings 10*, 450–463.
- Samuel, A. L. (1959). Some studies in machine learning using the game of checkers. *IBM Journal of research and development*, 3 (3), 210–229. <https://doi.org/10.1147/rd.33.0210>
- Schlenker, W., & Roberts, M. J. (2009). Nonlinear temperature effects indicate severe damages to US crop yields under climate change. *Proceedings of the National Academy of sciences*, 106 (37), 15594–15598. <https://doi.org/10.1073/pnas.0906865106>
- Schneider, U. A., Havlík, P., Schmid, E., Valin, H., Mosnier, A., Obersteiner, M., Böttcher, H., Skalský, R., Balkovič, J., & Sauer, T. (2011). Impacts of population growth, economic development, and technical change on global food production and consumption. *Agricultural Systems*, 104 (2), 204–215. <https://doi.org/10.1016/j.agsy.2010.11.003>
- Schultz, T. W. (1956). Reflections on agricultural production, output and supply. *Journal of farm Economics*, 38 (3), 748–762. <https://doi.org/10.2307/1234459>
- Singh, P. K., Pandey, A. K., Ahuja, S., & Kiran, R. (2022). Multiple forecasting approach: A prediction of co2 emission from the paddy crop in india. *Environmental Science and Pollution Research*, 1–12.

- Singh, S. R. (2007). A simple method of forecasting based on fuzzy time series. *Applied mathematics and computation*, 186 (1), 330–339. <https://doi.org/10.1016/j.amc.2006.07.128>
- Stas, M., Van Orshoven, J., Dong, Q., Heremans, S., & Zhang, B. (2016). A comparison of machine learning algorithms for regional wheat yield prediction using NDVI time series of SPOT-VGT. *2016 fifth international conference on agro-geoinformatics (agro-geoinformatics)*, 1–5.
- Su, Y.-x., Xu, H., & Yan, L.-j. (2017). Support vector machine-based open crop model (sbocm): Case of rice production in china. *Saudi journal of biological sciences*, 24 (3), 537–547.
- Troy, T. J., Kipgen, C., & Pal, I. (2015). The impact of climate extremes and irrigation on US crop yields. *Environmental Research Letters*, 10 (5), 054013. <https://doi.org/10.1088/1748-9326/10/5/054013>
- Van Klompenburg, T., Kassahun, A., & Catal, C. (2020). Crop yield prediction using machine learning: A systematic literature review. *Computers and Electronics in Agriculture*, 177, 105709. <https://doi.org/10.1016/j.compag.2020.105709>
- Vapnik, V. N. (1963). Pattern recognition using generalized portrait method. *Automation and remote control*, 24 (6), 774–780.
- Wang, Y., Zhang, Z., Feng, L., Du, Q., & Runge, T. (2020). Combining multi-source data and machine learning approaches to predict winter wheat yield in the conterminous united states. *Remote Sensing*, 12 (8), 1232. <https://doi.org/10.3390/rs12081232>
- Weber, J. G., Wall, C., Brown, J., & Hertz, T. (2015). Crop prices, agricultural revenues, and the rural economy¹. *Applied Economic Perspectives and Policy*, 37 (3), 459–476. <https://doi.org/10.1093/aep/ppy040>
- Wei, H.-y., Chen, Z.-f., Xing, Z.-p., Lei, Z., Liu, Q.-y., Zhang, Z.-z., Jiang, Y., HU, Y.-j., ZHU, J.-y., & CUI, P.-y. (2018). Effects of slow or controlled release fertilizer types and fertilization modes on yield and quality of rice. *Journal of Integrative Agriculture*, 17 (10), 2222–2234. [https://doi.org/10.1016/S2095-3119\(18\)62052-0](https://doi.org/10.1016/S2095-3119(18)62052-0)

- Wong, B. K., & Selvi, Y. (1998). Neural network applications in finance: A review and analysis of literature (1990–1996). *Information & management*, 34 (3), 129–139. [https://doi.org/10.1016/S0378-7206\(98\)00050-0](https://doi.org/10.1016/S0378-7206(98)00050-0)
- World Bank. (2023). *World Bank Agriculture Data* (tech. rep.) (Accessed on 16th December 2023). <https://data.worldbank.org/indicator/NV.AGR.TOTL.ZS?locations=NP>
- Yitayew, A., Abdulai, A., & Yigezu, Y. A. (2022). Improved agricultural input delivery systems for enhancing technology adoption: evidence from a field experiment in Ethiopia. *European Review of Agricultural Economics*, 49 (3), 527–556. <https://doi.org/10.1093/erae/jbac006>
- Yoo, C. K., Gopinath, M., & Kim, H. (2012). Trade policy reform, productivity growth and welfare in South Korean agriculture. *Applied Economic Perspectives and Policy*, 34 (3), 472–488.
- Yost, R., Uehara, G., Wade, M., Sudjadi, M., Widjaja-Adhi, I., & Zhi-Cheng, L. (1988). Expert systems in agriculture: Determining lime recommendations for soils of the humid tropics.